

eo-tides: Tide modelling tools for large-scale satellite Earth observation analysis

Robbi Bishop-Taylor¹, Claire Phillips¹, Stephen Sagar¹, Vanessa Newey¹, and Tyler Sutterley²

¹ Geoscience Australia, Australia ² University of Washington Applied Physics Laboratory, United States of America Corresponding author

DOI: [10.xxxxxx/draft](https://doi.org/10.xxxxxx/draft)

Software

- [Review](#)
- [Repository](#)
- [Archive](#)

Editor: [Open Journals](#)

Reviewers:

- [@openjournals](#)

Submitted: 01 January 1970

Published: unpublished

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

The eo-tides package provides powerful parallelised tools for integrating satellite Earth observation (EO) data with ocean tide modelling. The package provides a flexible Python toolkit for attributing modelled tide heights to a time-series of satellite images based on the spatial extent and acquisition time of each satellite observation (Figure 1).

eo-tides leverages advanced tide modelling functionality from the pyTMD tide prediction software (Sutterley et al., 2017), combining this capability with EO spatial analysis tools from the Open Data Cube (ODC)'s odc-geo (odc-geo contributors, 2024). This allows tides to be modelled in parallel using over 50 supported models, and returned in standardised pandas (McKinney, 2010; pandas development team, 2020) and xarray (Hoyer & Joseph, 2017) data formats for EO analysis.

eo-tides tools can be applied to petabytes of freely available satellite data loaded from the cloud using ODC's odc-stac or datacube packages (e.g. using Digital Earth Australia or Microsoft Planetary Computer's STAC SpatioTemporal Asset Catalogues). Additional functionality allows users to assess potential satellite-tide biases and validate modelled tides with external tide gauge data — critical considerations for ensuring the reliability and accuracy of coastal EO workflows. These open-source tools support the efficient, scalable and robust analysis of coastal EO data for any time period or location globally where ocean tide model data exists.




Figure 1: A typical eo-tides coastal EO workflow, with tide heights modelled into every pixel in a spatio-temporal stack of satellite data (e.g. Sentinel-2 or Landsat), then combined to derive insights into dynamic coastal environments.

Statement of need

Satellite remote sensing offers an unparalleled resource for examining dynamic coastal environments through time or across large regions (Turner et al., 2021; Vitousek et al., 2023). However, the highly variable influence of ocean tides can complicate analyses, making it difficult to separate the influence of changing tides from patterns of true coastal change (Vos et al., 2023). This is a particularly challenging for large-scale coastal EO analyses, where failing to account for tide dynamics can lead to inaccurate or misleading insights into satellite-observed coastal processes.

Conversely, information about ocean tides can provide unique environmental insights that can significantly enhance the value of EO data. Traditionally, satellite data dimensions include the geographic “where” and temporal “when” of acquisition. Introducing tide height as an additional analysis dimension allows data to be filtered, sorted, and analysed based on tidal dynamics, offering a transformative re-imagining of traditional multi-temporal EO analysis (Sagar et al., 2017). For instance, satellite data can be analysed to focus on ecologically significant tidal stages (e.g., high tide, low tide, spring or neap tides) or specific tidal processes (e.g., ebb or flow tides) (Sent et al., 2025).

This concept has been used to map coastal change at continental-scale (Bishop-Taylor et al., 2021), map intertidal zone extent and elevation (Bishop-Taylor et al., 2019; Fitton et al., 2021; Murray et al., 2012; Sagar et al., 2017), and creating tidally-constrained coastal image composites (Sagar et al., 2018). However, these methods have traditionally relied on bespoke, closed-source, or difficult-to-install tide modelling tools, limiting their reproducibility and portability. To support the next generation of coastal EO workflows, there is a pressing need for efficient open-source tools for combining satellite data with tide modelling. `eo-tides` addresses this need through functionality offered in five main analysis modules (`utils`, `model`, `eo`, `stats`, `validation`).

Features

Setting up tide models

The `eo_tides.utils` module simplifies the setup of ocean tide models, addressing a common barrier to coastal EO workflows. Tools like `list_models` provide feedback on available and supported models (Figure 2), while `clip_models` can significantly improve performance by clipping large high-resolution model files (e.g. FES2022 (Carrere et al., 2022)) to smaller study area extents.





	Model	Expected path
	EOT20	tide_models/EOT20/ocean_tides
	FES2022	tide_models/fes2022b/ocean_tide
	HAMTIDE11	tide_models/hamtide
...
Summary:		
Available models: 2/50		

Figure 2: A `list_tides` output providing a useful summary of available and supported tide models.

58 **Modelling tides**

59 The `eo_tides.model` module is powered by tide modelling functionality from the pyTMD Python
60 package (Sutterley et al., 2017). pyTMD is an open-source tidal prediction software that
61 simplifies the calculation of ocean and earth tides.

62 The `model_tides` function from `eo_tides.model` wraps pyTMD functionality to return tide
63 predictions in a standardised `pandas.DataFrame` format, enabling integration with EO data
64 and parallelisation for improved performance (Table 1). The `model_phases` function can
65 additionally classify tides into high/low/flow/ebb phases, critical for correctly interpreting
66 satellite-observed coastal processes like turbidity (Sent et al., 2025).

Table 1: A benchmark comparison of tide modelling parallelisation, for a typical large-scale analysis involving a month of hourly tides modelled at 10,000 points using three models (FES2022, TPXO10, GOT5.6).

Cores	Parallelisation	No parallelisation	Speedup
8	2min 46s ± 663 ms	9min 28s ± 536 ms	3.4x
32	55.9 s ± 560 ms	9min 24s ± 749 ms	10.1x

67 **Combining tides with satellite data**

68 The `eo_tides.eo` module integrates modelled tides with xarray-format satellite data (Hoyer &
69 Joseph, 2017). The `tag_tides` and `pixel_tides` functions (Table 2, Figure 3) can be applied
70 to attribute tides to satellite data for any coastal location on the planet, for example using
71 open data loaded from the cloud using ODC and STAC (STAC contributors, 2024).

Table 2: Comparison of the `tag_tides` and `pixel_tides` functions.

<code>tag_tides</code>	<code>pixel_tides</code>
<ul style="list-style-type: none">- Assigns a single tide height to each satellite image time-step- Single tide height per image can produce artefacts and discontinuities- Fast, low memory use- Ideal for small-scale analysis in non-complex tidal environments	<ul style="list-style-type: none">- Assigns a tide height to every individual pixel through time- Produce spatially seamless results across large regions- Slower, higher memory use- Ideal for large-scale analysis and coastal product generation




Figure 3: An example spatial tide height output produced by the `pixel_tides` function.

72 Calculating tide statistics and satellite biases

73 The `eo_tides.stats` module identifies biases caused by complex tide aliasing interactions that
74 can prevent satellites from observing the entire tide cycle (Bishop-Taylor et al., 2019; Eleveld
75 et al., 2014; Sent et al., 2025). The `tide_stats` and `pixel_stats` functions produce useful
76 statistics that summarise how well satellite data captures real-world tides (Figure 4).

🌊 Modelled astronomical tide range: 9.30 metres.
📡 Observed tide range: 6.29 metres.
● 68% of the modelled astronomical tide range was observed at this location.
● The highest 8% (0.77 metres) of the tide range was never observed.
● The lowest 24% (2.25 metres) of the tide range was never observed.
🌊 Mean modelled astronomical tide height: -0.00 metres.
📡 Mean observed tide height: 0.69 metres.




Figure 4: An example of tidally-biased satellite coverage, where only ~68% of the astronomical tide range is observed.

Validating modelled tides

The `eo_tides.validation` module validates modelled tides against observed sea-level measurements, assisting users to evaluate and select optimal models for their application (Figure 5).




Figure 5: A comparison of multiple tide models (EOT20, GOT5.5, HAMTIDE11) against observed sea level data from the Broome 62650 GESLA tide gauge (Haigh et al., 2023).

Research projects

Early versions of `eo-tides` functions have been used for continental-scale intertidal mapping (Bishop-Taylor et al., 2024), multi-decadal shoreline mapping across Australia (Bishop-Taylor et al., 2021) and Africa, and for correcting satellite-derived shoreline in the CoastSeg Python package (Fitzpatrick et al., 2024).

Acknowledgements

Functions from `eo-tides` were originally developed in the Digital Earth Australia Notebooks repository (Krause et al., 2021). This paper is published with the permission of the Chief Executive Officer, Geoscience Australia.

References

- Bishop-Taylor, R., Nanson, R., Sagar, S., & Lymburner, L. (2021). Mapping Australia's Dynamic Coastline at Mean Sea Level using Three Decades of Landsat Imagery. *Remote Sensing of Environment*, 267, 112734. <https://doi.org/10.1016/j.rse.2021.112734>
- Bishop-Taylor, R., Phillips, C., Newey, V., & Sagar, S. (2024). *Digital Earth Australia Intertidal*. Commonwealth of Australia (Geoscience Australia). <https://doi.org/10.26186/149403>
- Bishop-Taylor, R., Sagar, S., Lymburner, L., & Beaman, R. J. (2019). Between the tides: Modelling the elevation of australia's exposed intertidal zone at continental scale. *Estuarine, Coastal and Shelf Science*, 223, 115–128. <https://doi.org/10.1016/j.ecss.2019.03.006>
- Carrere, L., Lyard, F., Cancet, M., Allain, D., Dabat, M.-L., Fouchet, E., Sahuc, E., Faugere, Y., Dibarboure, G., & Picot, N. (2022). A new barotropic tide model for global ocean: FES2022. *2022 Ocean Surface Topography Science Team Meeting*, 43. <https://doi.org/10.24400/527896/a03-2022.3287>
- Eleveld, M. A., Van der Wal, D., & Van Kessel, T. (2014). Estuarine suspended particulate matter concentrations from sun-synchronous satellite remote sensing: Tidal and meteorological effects and biases. *Remote Sensing of Environment*, 143, 204–215. <https://doi.org/10.1016/j.rse.2013.12.019>

- 106 Fitton, J. M., Rennie, A. F., Hansom, J. D., & Muir, F. M. E. (2021). Remotely sensed
107 mapping of the intertidal zone: A Sentinel-2 and Google Earth Engine methodology. *Remote*
108 *Sensing Applications: Society and Environment*, 22(100499), 1–12. <https://doi.org/10.1016/j.rsase.2021.100499>
- 110 Fitzpatrick, S., Buscombe, D., Warrick, J. A., Lundine, M. A., & Vos, K. (2024). CoastSeg: An
111 accessible and extendable hub for satellite-derived-shoreline (SDS) detection and mapping.
112 *Journal of Open Source Software*, 9(99), 6683. <https://doi.org/10.21105/joss.06683>
- 113 Haigh, I. D., Marcos, M., Talke, S. A., Woodworth, P. L., Hunter, J. R., Hague, B. S.,
114 Arns, A., Bradshaw, E., & Thompson, P. (2023). GESLA version 3: A major update to
115 the global higher-frequency sea-level dataset. *Geoscience Data Journal*, 10(3), 293–314.
116 <https://doi.org/10.1002/gdj3.174>
- 117 Hoyer, S., & Joseph, H. (2017). xarray: N-d labeled arrays and datasets in python. *Journal of*
118 *Open Research Software*, 5(1). <https://doi.org/10.5334/jors.148>
- 119 Krause, C., Dunn, B., Bishop-Taylor, R., Adams, C., Burton, C., Alger, M., Chua, S., Phillips, C.,
120 Newey, V., Kouzoubov, K., Leith, A., Ayers, D., & Hicks, A. (2021). *Digital Earth Australia*
121 *notebooks and tools repository*. <https://github.com/GeoscienceAustralia/dea-notebooks/>;
122 Commonwealth of Australia (Geoscience Australia). <https://doi.org/10.26186/145234>
- 123 McKinney, Wes. (2010). Data Structures for Statistical Computing in Python. In Stéfan van
124 der Walt & Jarrod Millman (Eds.), *Proceedings of the 9th Python in Science Conference*
125 (pp. 56–61). <https://doi.org/10.25080/Majora-92bf1922-00a>
- 126 Murray, N. J., Phinn, S. R., Clemens, R. S., Roelfsema, C. M., & Fuller, R. A. (2012).
127 Continental scale mapping of tidal flats across east asia using the landsat archive. *Remote*
128 *Sensing*, 4(11), 3417–3426. <https://doi.org/10.3390/rs4113417>
- 129 odc-geo contributors. (2024). Opendatacube/odc-geo. In *GitHub repository*. GitHub.
130 <https://github.com/opendatacube/odc-geo>
- 131 pandas development team. (2020). *Pandas-dev/pandas: pandas* (latest). Zenodo. <https://doi.org/10.5281/zenodo.3509134>
- 133 Sagar, S., Phillips, C., Bala, B., Roberts, D., Lymburner, L., & Beaman, R. J. (2018).
134 Generating continental scale pixel-based surface reflectance composites in coastal regions
135 with the use of a multi-resolution tidal model. *Remote Sensing*, 10(3), 480. <https://doi.org/10.3390/rs10030480>
- 137 Sagar, S., Roberts, D., Bala, B., & Lymburner, L. (2017). Extracting the intertidal extent and
138 topography of the australian coastline from a 28 year time series of landsat observations.
139 *Remote Sensing of Environment*, 195, 153–169. <https://doi.org/10.1016/j.rse.2017.04.009>
- 140 Sent, G., Antunes, C., Spyrakos, E., Jackson, T., Atwood, E. C., & Brito, A. C. (2025). What
141 time is the tide? The importance of tides for ocean colour applications to estuaries. *Remote*
142 *Sensing Applications: Society and Environment*, 37, 101425. <https://doi.org/10.2139/ssrn.4858713>
- 144 STAC contributors. (2024). *SpatioTemporal Asset Catalog (STAC) specification*. <https://stacspec.org>
- 146 Sutterley, T. C., Alley, K., Brunt, K., Howard, S., Padman, L., & Siegfried, M. (2017). *pyTMD:*
147 *Python-based tidal prediction software*. Zenodo. <https://doi.org/10.5281/zenodo.5555395>
- 148 Turner, I. L., Harley, M. D., Almar, R., & Bergsma, E. W. J. (2021). Satellite optical imagery
149 in Coastal Engineering. *Coastal Engineering*, 167, 103919. <https://doi.org/10.1016/j.coastaleng.2021.103919>
- 151 Vitousek, S., Buscombe, D., Vos, K., Barnard, P. L., Ritchie, A. C., & Warrick, J. A. (2023).
152 The future of coastal monitoring through satellite remote sensing. *Cambridge Prisms:*

153 *Coastal Futures*, 1, e10. <https://doi.org/10.1017/cft.2022.4>

154 Vos, K., Splinter, K. D., Palomar-Vázquez, J. E., Almonacid-Caballer, J., Cabezas-Rabadán,
155 C., Kras, E. C., Luijendijk, A. P., Calkoen, F., Almeida, L. P., Pais, D., Klein, A. H. F.,
156 Mao, Y., Harris, D., Castelle, B., Buscombe, D., & Vitousek, S. (2023). Benchmarking
157 satellite-derived shoreline mapping algorithms. *Communications Earth & Environment*,
158 4(345). <https://doi.org/10.1038/s43247-023-01001-2>

DRAFT