

EXOplanet Transit Interpretation Code (EXOTIC) Instructions: How to Run the Code

I. Using the Terminal/Ubuntu/ Windows CMD Terminal (skip this step if you are already familiar with the command line)


- Opening the Application
 - If you are using a Mac, open up the terminal.
 - If you followed the recommended installation instructions for Windows and are using Ubuntu, open Ubuntu.
 - If you chose to run EXOTIC natively on Windows, open Command Prompt.
- Changing Directories
 - A “directory” is just another term for a folder in your computer
 - To enter a directory in terminal, type “cd” followed by a space, and then the path to your directory, and then hitting return which will execute the command.
 - The path to the folder you want to change into can be found by right clicking on the folder in Finder, and then holding down the option key, and then selecting “Copy ‘Your file’ as Pathname”
 - Pasting that copied file path after the “cd” is the easiest way to accurately and reliably change into your desired directory
 - To go back to your “Home” directory, simply type “cd”, or “cd ~” for Windows CMD users and then hit enter
 - “pwd” (“cd” for those using Windows CMD Terminal) can be used to determine your current directory
- Listing Files in a Directory
 - To list all of the files in your current directory, enter “ls”, or “dir” for those using Windows CMD Terminal)
 - This is a good way to make sure that a/e file you are looking for is actually in the directory you think it is
- Keyboard Interrupt
 - To kill a process or program while its running, hit “Ctrl + c”
- Quitting the Terminal
 - To quit the terminal, after you have killed any processes that were running using the keyboard interrupt, type exit and hit enter



II. Getting the Code to Run

If you are using a Mac or Ubuntu:

1. Change into the directory in which “exotic.py” is located using Terminal (`cd .../EXOTIC`)
2. Type “`python3 exotic.py`” and then hit enter
 - This command may be “`python exotic.py`”. Run this instead if command is not found.
3. If you see the EXOTIC header, the pipeline is running!



```
mjs2369@DESKTOP-R2PG3MN: /mnt/c/Users/Marlena Smith/Desktop/ExoplanetWatch/EXOTIC
mjs2369@DESKTOP-R2PG3MN:/mnt/c/Users/Marlena Smith/Desktop/ExoplanetWatch/EXOTIC$ cd /mnt/c/Users/Marlena Smith/Desktop/ExoplanetWatch/EXOTIC
mjs2369@DESKTOP-R2PG3MN:/mnt/c/Users/Marlena Smith/Desktop/ExoplanetWatch/EXOTIC$ python3 exotic.py
Importing Python Packages - please wait.
Thinking -WARNING (theano.tensor.blas): Using NumPy C-API based implementation for BLAS functions.
Thinking -


*****
Welcome to the EXOplanet Transit Interpretation Code (EXOTIC)
Version 0.7.5
*****

Enter "1" for Real Time Reduction or "2" for for Complete Reduction:
```

If you are using Windows CMD Terminal:

1. Change into the directory in which “exotic.py” is located using CMD Terminal
2. Type “`ipython`” and then hit enter
3. Type “`run exotic.py`” and then hit enter

4. If you see the header for EXOTIC, the pipeline is running!



```
IPython: C:\ExoplanetWatch\EXOTIC
C:\Users\Marlena Smith\Desktop\ExoplanetWatch>cd EXOTIC
C:\Users\Marlena Smith\Desktop\ExoplanetWatch\EXOTIC>ipython
Python 3.7.6 (default, Jan 8 2020, 20:23:39) [MSC v.1916 64 bit (AMD64)]
Type 'copyright', 'credits' or 'license' for more information
IPython 7.12.0 -- An enhanced Interactive Python. Type '?' for help.

In [1]: run exotic.py
Importing Python Packages - please wait.
Thinking /gzip was not found on your system! You should solve this issue for astroquery.eso to be at its best!
On POSIX system: make sure gzip is installed and in your path!On Windows: same for 7-zip (http://www.7-zip.org)!
Thinking /WARNING (theano.configdefaults): g++ not available, if using conda: `conda install m2w64-toolchain`
C:\anaconda3\lib\site-packages\theano\configdefaults.py:560: UserWarning: DeprecationWarning: there is no c++ compiler. This is deprecated and with Theano 0.11 a c++ compiler will be mandatory
  warnings.warn("DeprecationWarning: there is no c++ compiler.")
WARNING (theano.configdefaults): g++ not detected ! Theano will be unable to execute optimized C-implementations (for both CPU and GPU) and will default to Python implementations. Performance will be severely degraded. To remove this warning, set Theano flags cxx to an empty string.
Thinking /WARNING (theano.tensor.blas): Using NumPy C-API based implementation for BLAS functions.
Thinking -

*****
Welcome to the EXoplanet Transit Interpretation Code (EXOTIC)
Version 0.7.5
*****

Enter "1" for Real Time Reduction or "2" for Complete Reduction:
```

After the program is running, you will first be presented with 2 options:

Enter '1' to Reduce your Data in Real Time

- i. This option should only be selected if you want to a quick reduction your data while you are on an observing run
- ii. This will simply serve as a visual tool, but will not fit a model lightcurve, nor will it extract planetary data

Enter '2' to do a Complete Reduction


- iii. The default option that should be selected at all times except for the case described in 1ai.
- iv. This performs a full reduction on all of your data
 1. See How it Works for an explanation of the reduction process

III. Real Time Reduction Routine (skip this if you entered “2”)

Now that you have the pipeline running, this chapter will walk you through how to reduce your dataset in real time. The photos used below are using an example data set of HAT-P-32b.

1. Enter the directory path to the FITS Image Files
 - a. You will need to enter the complete path to the folder where the images you will be taking will get saved into

- i. If you don't have a folder in which your images will get saved, you will need to make one by simply creating a new folder using finder
 - b. The easiest way to get the path to your folder is by right clicking on the folder in Finder, and then holding down the option key, selecting "Copy 'Your file' as Pathname", and then pasting that into the terminal
2. Enter the name of the planet you are observing
 - a. For plot title only
3. Start Your Observing Run
 - a. Once you start your observing run, ".FITS" files should be entering the directory at the rate you set your observing cadence to.
4. Type "continue" after the first image has been taken and saved after the prompt
5. Entering the Pixel Coordinates of your Target Star
 - a. Go to: <http://nova.astrometry.net/upload>
 - b. Upload the first ".FITS" image in your in your dataset to Astrometry.net (left)
 - c. After it finishes analyzing your image click "Go to results page" (center)
 - d. Then click to download "new-image.fits" (right)



- e. Open "new-image.fits" in DS9, which should now be WCS encoded
 - i. To open a fits image with DS9, simply right click on the image in Finder, and select "Open With" and choose "SAOImageDS9"
 - ii. When opening for the first time, the system will say the application is by an unidentified developer. To get past this, open System Preferences and under the General pane of the Security & Privacy tab, you can allow the application under "Allow apps downloaded from:"
 - iii. Here is the User Guide: [DS9 User Guide](#) on their website
 - f. Search the SIMBAD database for your target star by typing its name into the search box
 - i. Link: [SIMBAD Search Portal](#)

- ii. Once you see a page like this one, with your target name, obtain the RA and Dec Coordinates (Green Box) and note the V-mag (Orange Box)

hat p 32

other query modes : [Identifier query](#) [Coordinate query](#) [Criteria query](#) [Reference query](#) [Basic query](#) [Script submission](#) [TAP](#) [Output options](#) [Help](#)

Query : hat p 32

C.D.S. - SIMBAD4 rel 1.7 - 2019.07.04CEST01:56:07

Available data : [Basic data](#) • [Identifiers](#) • [Plot & images](#) • [Bibliography](#) • [Measurements](#) • [External archives](#) • [Notes](#) • [Annotations](#)

Basic data :

HAT-P-32 -- Star

Other object types:

ICRS coord. (*ep*=J2000) : 02 04 10.2775457769 +46 41 16.210382751 Optical) [0.0572 0.0399 90] A 2018yCat.1345....0G

FK4 coord. (*ep*=B1950 *eq*=1950) : 02 04 10.2775457769 +46 41 16.210382751 [0.0572 0.0399 90]

Gal coord. (*ep*=J2000) : 135.6985495038660 -14.3730606007401 [0.0572 0.0399 90]

Proper motions *mas/yr* : -9.825 3.477 [0.100 0.086 90] A 2018yCat.1345....0G

Radial velocity / Redshift / *cz* : V(km/s) -23.95 [0.23] / *z*(spectroscopic) -0.000080 [0.000001] / *cz* -23.95 [0.23] (Opt) A 2018yCat.1345....0G

Parallaxes (*mas*): 3.4305 [0.0624] A 2018yCat.1345....0G

Fluxes (7) : V 11.44 [0.12] D 2000AA...355L...27H
 R 11.23 [0.06] E 2003yCat.1315....0Z
 G 11.1335 [0.0005] C 2018yCat.1345....0G
 J 10.251 [0.022] C 2003yCat.2246....0C
 H 10.024 [0.022] C 2003yCat.2246....0C
 K 9.990 [0.022] C 2003yCat.2246....0C


SIMBAD with radius arcmin

Interactive AladinLite view


- g. Find your target on the “new-image.fits” in DS9 by moving your cursor around on the image until the “fk5” values match the Ra and Dec values shown on the SIMBAD Database
- i. You can see stars more clearly by clicking “scale” → “zscale”

- h. When the FK5 values match the Ra and Dec from Simbad, the numbers in the boxes next to image (blue box) are the x and y pixel coordinates of your target

Move your cursor around the image (right) until your RA and Dec shown on the green box on the image match the values on SIMBAD



6. Entering the Pixel Coordinates of your Comparison Star
 - a. Return to the SIMBAD Page of your Star
 - b. Above the image of your target on SIMBAD, click on the “Interactive AladinLite View”
 - c. Check the box that says SIMBAD under “Catalogues”
 - d. Then click on the nearby stars (green circles) around your target star (purple +), looking for stars of similar brightness
 - e. Then in the “SIMBAD source” box on the left, check to make sure you have in fact clicked on a star by ensuring “main_type” = Star
 - f. Then scroll down in the “SIMBAD source” box and check the number next to V. If it is close to the V-mag of your target star (within 1.5 units), you have likely found a good comparison star!
 - g. Obtain the “ra_sex” and “dec_sex” which are located in the “SIMBAD source” box (green arrow)
 - i. These are the RA and Dec values for your comp star




- h. Find your comparison star on your “new-image.fits” in DS9 using the same procedure you did for finding your target star. (step 4h)
 - i. Enter the X and Y coordinates of your comparison star into your terminal.
7. A plot of Normalized Flux vs. Time (a lightcurve!) will be generated, and this plot will get updated every time a new image is added to the specified folder.
8. After you are done observing, either close out the plot, or quit the program by typing “Ctrl+c”

IV. Complete Reduction Routine

Now that you have the pipeline running, this chapter will walk you through how to perform a full reduction of your dataset. The photos used below are using an example data set of HAT-P-32b, which can be found in the sample-data folder.

1. You will first be asked whether you will be using .FITS files or pre-reduced data in a .txt file.
 - a. If you are unsure of which one you are using, you will most likely be using .FITS files (these files contain raw data obtained directly from your telescope).
 - b. Type the correct number and press enter.
2. You will then be asked how you would like to input your initial parameters.
 - a. Enter “1” to do so manually and **follow steps 3-10 below**.
 - b. Enter “2” to do so using an input file, which is a file that contains all the parameters EXOTIC will ask for, including the location of your data.
 - i. Enter the directory and filename of the input file. Hit enter.
 - ii. If you select this option, **skip to step 11 below**.
 - iii. You can test this method by using the file in your EXOTIC folder titled “inits.txt”. This input file contains the figures for the star HAT-P-32b, which corresponds to the sample data.

- iv. To learn more about input files and how to create and utilize them, check out our guide on Input Files:
<https://github.com/rzellem/EXOTIC/tree/main/Documentation>
3. Enter the directory path to the FITS Image Files
 - a. You will need to enter the complete path to the folder where the images you want to reduce are located
 - b. The easiest way to get the path to your folder is by right clicking on the folder in Finder, and then holding down the option key, selecting “Copy ‘Your file’ as Pathname”, and then pasting that into the terminal
4. Enter the directory path to the folder you want the results of the reduction saved into
 - i. If you don’t already have a folder where you want your plots saved into, you will need to make one by simply creating a new folder using finder in your desired location
5. Enter the name of the planet you are looking at and then the date (MM-DD-YYYY)
 - a. This information is just used to make plot titles. It is not relevant to the data reducing itself.
6. Enter the longitude and latitude in degrees of where you observed when prompted
 - a. These values are easy to find by simply googling it (an estimate is sufficient)
7. Locate Your Target Star
 - a. Go to: <http://nova.astrometry.net/upload>
 - b. Upload the first “.FITS” image in your in your dataset to Astrometry.net (left)
 - c. After it finishes analyzing your image click “Go to results page” (center)
 - i. If the results shown in the figures do not appear, refresh the page.
 - d. Then click to download “new-image.fits” (right)



- ii. Use the name formatting from the source tweet
- g. Once you see a page like this one, with your target name, obtain the RA and Dec Coordinates (Green Box) and note the V-mag (Orange Box)

hat p 32

other query modes :

Identifier query

Coordinate query

Criteria query

Reference query

Basic query

Script submission

TAP

Output options

Help

Query : hat p 32

C.D.S. - SIMBAD4 rel 1.7 - 2019.07.04CEST01:56:07

Available data : [Basic data](#) • [Identifiers](#) • [Plot & images](#) • [Bibliography](#) • [Measurements](#) • [External archives](#) • [Notes](#) • [Annotations](#)

Basic data :

HAT-P-32 -- Star

Other object types:

ICRS coord. (*ep*=J2000) : 02 04 10.2775457769 +46 41 16.210382751 Optical) [0.0572 0.0399 90] A 2018yCat.1345....0G

FK4 coord. (*ep*=B1950 *eq*=1950) : 02 01 00.5974025128 +46 26 53.987833965 [0.0572 0.0399 90]

Gal coord. (*ep*=J2000) : 135.6985495038660 -14.3730606007401 [0.0572 0.0399 90]

Proper motions *mas/yr* : -9.825 3.477 [0.100 0.086 90] A 2018yCat.1345....0G

Radial velocity / Redshift / *cz* : V(km/s) -23.95 [0.23] / *z*(spectroscopic) -0.000080 [0.000001] / *cz* -23.95 [0.23] (Opt) A 2018yCat.1345....0G

Parallaxes (*mas*): 3.4305 [0.0624] A 2018yCat.1345....0G

Fluxes (7) : B 11.79 [0.10] D 2000A&A...355L...27H
V 11.44 [0.12] D 2000A&A...355L...27H
R 11.23 [0.06] E 2009yCat.1315....0Z
G 11.1335 [0.0005] C 2018yCat.1345....0G
J 10.251 [0.022] C 2003yCat.2246....0C
H 10.024 [0.022] C 2003yCat.2246....0C
K 9.990 [0.022] C 2003yCat.2246....0C


SIMBAD with radius arcmin




- h. Enter the Ra and Dec in the form ‘##:##:##’
 - i. For example, in the example above, the Ra should get entered as 02:04:10 and the Dec as 46:41:16 (ignore the values after the decimal)
- i. Find your target on the “new-image.fits” in DS9 by moving your cursor around on the image until the “fk5” values match the Ra and Dec values shown on the SIMBAD Database
 - i. You can see stars more clearly by clicking “scale” → “zscale”
 1. Blue highlighted buttons in image below

- ii. When the FK5 values match the Ra and Dec from Simbad, the numbers in the boxes next to image (blue box) are the x and y pixel coordinates of your target

Move your cursor around the image (right) until your RA and Dec shown on the green box on the image match the values on SIMBAD



- j. Enter the Target X and Y Image (pixel) position (Blue Box in image above) into the terminal when prompted
8. Decide how many comparison stars you want to use
 - a. 3 is recommended, but any number between 1-10 will do
9. Entering the X and Y Pixel Coordinates of your Comparison Stars
 - a. Return to the SIMBAD Page of your Star
 - b. Above the image of your target on SIMBAD, click on the “Interactive AladinLite View”
 - c. Check the box that says SIMBAD under “Catalogues”
 - d. Then click on the nearby stars (green circles) around your target star (purple +), looking for stars of similar brightness
 - e. Then in the “SIMBAD source” box on the left, check to make sure you have in fact clicked on a star by ensuring “main_type” = Star
 - f. Then scroll down in the “SIMBAD source” box and check the number next to V. If it is close to the V-mag of your target star, you have likely found a good comparison star!
 - g. Obtain the “ra_sesa” and “dec_sesa” which are located in the “SIMBAD source” box (green arrow)
 - i. These are the RA and Dec values for your comp star



- h. Find your comparison star on your “new-image.fits” in DS9 using the same procedure you did for finding your target star. (step 4h)
- i. Enter the X and Y coordinates of your comparison star into your terminal.
- j. Repeat steps 5b through 5j until you have entered the coordinates for each of the comparison stars you wanted to use
10. When asked if you have calibration images (flats, darks, or biases), enter ‘y’ if you have them, otherwise enter ‘n’
 - a. If you don’t know, or aren’t sure what they are, you probably don’t have them and should enter ‘n’ and the code will take you to step 8.
 - b. For each type of calibration image you have, enter the directory path to where each set located when prompted
11. Enter the Planetary Parameters for Lightcurve Fitting
 - a. Go to the NASA Exoplanet Archive look up
 - i. Link: [NASA Exoplanet Archive Search](https://exoplanetarchive.nasa.gov/)
 - b. Enter the name of your target star in the search bar under “Host Name” then click on its name once you have found it
 - i. The site is case sensitive so be careful with capitalization
 - ii. It should open a site that looks like the image shown at the bottom of page 11
 - c. The code will iterate through each parameter it needs and will attempt to scrape the values off of the Exoplanet Archive website
 - d. It will then prompt you to verify that each parameter is correct
 - i. Parameters: Orbital Period, Mid-Transit Time, Eccentricity, Ratio of Planet to Stellar Radius, Ratio of Distance to Stellar Radius, Orbital Inclination (all boxed in green in the image below)
 - ii. For Mid-Transit Time, you want to pick the largest value in the column and its corresponding uncertainty (the values after the \pm).

- iii. For everything else, pick the value highlighted in gold if it's available, and then go to the most recently published non-null value if it's not
- e. For each parameter it asks if you agree for, if the value the code obtained is not the same as what you see on the Archive, type 'n', and then enter the correct value when prompted

NASA Exoplanet Archive Links					
Planet	Related Overviews				Transit Service
	Confirmed		Kepler Pipeline		
HAT-P-32 b	Planet	Host			HAT-P-32 b Transits

Planet Orbital Properties								
Planet	Period (days)	Semi-Major Axis (AU)	Inclination (deg)	Eccentricity	Time of Periastron Passage (days)	Longitude of Periastron (deg)	Date of Orbital Solution	Reference
b	2.15000825±0.00000012	0.03427 ^{+0.00040} _{-0.00042}	88.90±0.40	<0.044	null	null	null	Bonomo et al. 2017
b	2.150010±0.000001	null	88.90±0.40	0.0	null	null	null	Stassun et al. 2017
b	2.1500080	null	89.33 ^{+0.58} _{-0.80}	0	null	null	null	Nortmann et al. 2016
b	2.15000805 ^{+0.00000093} _{-0.00000097}	0.0343±0.0004	88.9±0.4	0.0072 ^{+0.0700} _{-0.0064}	null	96 ^{+180.0} ₋₁₁	null	Zhao et al. 2014
b	null	null	null	0.20 ^{+0.19} _{-0.13}	null	58.0 ^{+28.0} _{-53.0}	null	Knutson et al. 2014
b	2.1500085±0.0000002	null	89.12 ^{+0.61} _{-0.68}	null	null	null	null	Gibson et al. 2013
b	2.150008±0.000001	0.0343±0.0004	88.9±0.4	0.	null	0	null	Hartman et al. 2011

Planet Parameters										
Planet	M sin(i)		Mass		Radius			Density	Equilibrium Temperature	Reference
	(Jupiter Mass)	(Earth Mass)	(Jupiter Mass)	(Earth Mass)	(Solar Radii)	(Jupiter Radii)	(Earth Radii)	(g/cm ³)	(K)	
b	null	null	0.75±0.13	238±41	0.1838±0.0026	1.789±0.025	20.05±0.28	0.163 ^{+0.029} _{-0.028}	null	Bonomo et al. 2017
b	null	null	0.830±0.210	264±67	0.180±0.014	1.75±0.14	19.6±1.6	null	null	Stassun et al. 2017
b	null	null	null	null	null	null	null	null	null	Nortmann et al. 2016
b	null	null	null	null	0.184±0.003	1.789±0.025	20.053±0.280	null	null	Zhao et al. 2014
b	null	null	0.79±0.15	251±48	null	null	null	null	null	Knutson et al. 2014
b	null	null	0.8600±0.1640	273.3±52.1	0.1845 ^{+0.0029} _{-0.0028}	1.796 ^{+0.028} _{-0.027}	20.13 ^{+0.31} _{-0.30}	0.18±0.04	1779±26	Gibson et al. 2013
b	null	null	0.860±0.164	273.322±52.122	0.1839±0.0026	1.789±0.025	20.053±0.280	0.19±0.04	1786±26	Hartman et al. 2011

Planet Transit Properties									
Planet	Depth (perc)	Duration (days)	Duration (hours)	Mid-Point (days)	Impact Parameter	Occultation Depth (perc)	Ratio of Distance to Stellar Radius	Ratio of Planet to Stellar Radius	Reference
b	null	null	null	2454420.44645±0.00009	null	null	null	null	Bonomo et al. 2017
b	2.300±0.012	null	null	null	null	null	6.06±0.13	null	Stassun et al. 2017
b	null	null	null	2456185.602987±0.000110	null	null	6.123 ^{+0.021} _{-0.054}	0.1516376 ^{+0.0008740} _{-0.0005450}	Nortmann et al. 2016
b	null	null	null	2454420.44712 ^{+0.000092} _{-0.000084}	null	null	null	null	Zhao et al. 2014
b	null	null	null	null	null	null	null	null	Knutson et al. 2014
b	null	0.12959 ^{+0.00059} _{-0.00057}	3.1102 ^{+0.0142} _{-0.0137}	2454942.898449±0.000077	0.0930 ^{+0.0710} _{-0.0650}	null	6.091 ^{+0.036} _{-0.047}	0.1515±0.0012	Gibson et al. 2013
b	1.8523	0.1295±0.0003	3.1080±0.0072	2454420.44637±0.00009	0.117 ^{+0.045} _{-0.047}	null	6.05 ^{+0.03} _{-0.04}	0.1508±0.0004	Hartman et al. 2011

12. Limb Darkening

- a. On the NASA Exoplanet Archive page of your planet, scroll down until you reach the section called “Summary of Stellar Information”

Summary of Stellar Information			
Right Ascension	02h04m10.28s	Declination	+46d41m16.2s
Galactic Longitude (deg)	135.69859	Galactic Latitude (deg)	-14.37298
Parallax (mas)	3.4304647653712994±0.0623970254391056	Distance (pc)	291.50569045176132±5.30222265110304
RA Proper Motion (mas/yr)	-9.8±0.1	Dec Proper Motion (mas/yr)	3.5±0.1
Total Proper Motion (mas/yr)	10.4±0.1	Radial Velocity (km/s)	-23.21±0.26
B-band (mag)	null	K-band (mag)	9.990±0.022
Spectral Type	null	Effective Temperature (K)	6207±88
Surface Gravity (log ₁₀ (cm/s ²))	4.33±0.01	Luminosity (log ₁₀ (L _{sun}))	null
Radius (R _{sun})	1.19±0.10	Mass (M _{sun})	1.09±0.28
Density (g/cm ³)	0.91±0.06	Age (Gyr)	null
Metallicity (dex)	-0.04	Metallicity Ratio	[Fe/H]
V sin(i) (km/s)	20.7±0.5	S-index	null
log R' ^{HK}	null	X-ray activity, log(L _x)	null
Number of Hipparcos Light Curves	0	Number of Photometric non-Hipparcos Light Curves	0
Number of Radial Velocity Time Series	0	Number of Amateur Light Curves	0
Number of Spectra	0	Number of Images	3

- i. It will ask for the Effective Temperature (green box), the Metallicity (red), and the Surface Gravity in the log(g) box (yellow)
13. Now that you have entered all the necessary information, the code will then reduce the dataset
14. After the reduction component is done a set of plots are generated and saved into your specified directory
 - a. The target star’s raw time varying flux (see How it Works sect...)
 - i. Figure name: ‘TargetRawFlux<planet name><date>.png’
 - b. The comparison star’s raw time varying flux
 - i. Figure name: ‘CompRawFlux<planet name><date>.png’
 - c. The Normalized Target Flux as a function of phase
 - i. Figure name: ‘NormalizedFluxPhase<planet name><date>.png’
15. The code uses an MCMC to fit a model light curve to your data, fitting for airmass, mid transit time, and planet radius
 - a. Warning: this will take some time. Usually around 30 minutes
 - b. For explanation, see How it Works: Full Lightcurve Fitting Routine
16. Final Light Curve fitted with a model is generated and saved in your specified directory
 - a. Figure name: ‘FinalLightCurve<planet name><date>.pdf’
17. The final planetary parameters of Mid Transit Time and Ratio of Planet to Stellar Radius and their corresponding uncertainties will be displayed in the terminal, and also saved as a ‘.txt’ file in your specified directory
 - a. Figure name: ‘FinalParams<planet name><date>.txt’

- b. To learn how to interpret your results, go to “How it Works: Results Interpretation”
- 18. If you want to save your reduced data into the official AAVSO data format file, type ‘y’ when prompted
 - a. Enter your AAVSO Observer ID
 - i. If you don’t have an Observer ID, it’s probably because you haven’t created an account, which can be done at:
<https://www.aavso.org/apps/register/>
 - b. Enter your exposure length in seconds
 - i. This is simply how long you exposed for when you took the data
 - ii. It often can be found by opening one of the images in DS9, and then going to File -> Display Header
 - c. Enter the binning used on your camera
 - i. If you don’t know or are unsure, just type ‘1x1’
 - d. The file should now be saved as “AAVSO<target name><date>.txt”